

Award Number: W81XWH-13-1-0040

TITLE: Role of CTGF in White Matter Development in Tuberous Sclerosis

PRINCIPAL INVESTIGATOR: Mustafa Sahin

CONTRACTING ORGANIZATION: The Children's Hospital Corporation
Boston, MA 02115

REPORT DATE: February 2015

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

| Report Documentation Page | | | Form Approved OMB No. 0704-0188 | | |
|---|------------------------------------|---|--|---|------------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE FEB 2015 | | 2. REPORT TYPE | | 3. DATES COVERED 01-02-2014 to 31-01-2015 | |
| 4. TITLE AND SUBTITLE Role of CTGF in White Matter Development in Tuberous Sclerosis | | 5a. CONTRACT NUMBER | | | |
| | | 5b. GRANT NUMBER | | | |
| | | 5c. PROGRAM ELEMENT NUMBER | | | |
| 6. AUTHOR(S) | | 5d. PROJECT NUMBER | | | |
| | | 5e. TASK NUMBER | | | |
| | | 5f. WORK UNIT NUMBER | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Children's Hospital Corporation,,Boston,,MA,02115 | | 8. PERFORMING ORGANIZATION REPORT NUMBER | | | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | 10. SPONSOR/MONITOR'S ACRONYM(S) | | | |
| | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | | | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT Our results show that neurons lacking Tsc1 results in a block in oligodendrocyte maturation and thus hypomyelination in the mouse brain. We show that Tsc-deficient neurons secrete excessive amounts of connective tissue growth factor (CTGF), which blocks the maturation of oligodendrocytes. We show here that CTGF is both necessary and sufficient to block the oligodendrocyte maturation. To determine whether CTGF is necessary in vivo, we generated mice lacking both Tsc1 and CTGF in neurons (Tsc1cc;CTGFff/f;SynCre+) mice. In order to investigate the effect of CTGF on oligodendrocyte maturation, we propose to treat the oligodendrocytes with different domains of CTGF (Module I to IV). Moreover, in order to investigate the upstream regulation of CTGF expression in Tsc-deficient neurons, we tested Serum Response Factor (SRF) pathway, which was previously shown to affect CTGF expression. Our preliminary results show that SRF is downregulated in Tsc1 mutant brains and this can be rescued by rapamycin treatment suggesting a crosstalk between SRF and mTOR pathways. | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT Same as Report (SAR) | 18. NUMBER OF PAGES 11 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

| REPORT DOCUMENTATION PAGE | | | Form Approved OMB No. 0704-0188 | | |
|---|------------------|--------------------------|---|--|---|
| Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. | | | | | |
| 1. REPORT DATE february 2015 | | 2. REPORT TYPE Annual | | 3. DATES COVERED 1 Feb 2014 - 31 Jan 2015 | |
| 4. TITLE AND SUBTITLE Role of CTGF in White Matter Development in Tuberous Sclerosis | | | 5a. CONTRACT NUMBER | | |
| | | | 5b. GRANT NUMBER W81XWH-13-1-0040 | | |
| | | | 5c. PROGRAM ELEMENT NUMBER | | |
| 6. AUTHOR(S) Mustafa Sahin email: mustafa.sahin@childrens.harvard.edu | | | 5d. PROJECT NUMBER | | |
| | | | 5e. TASK NUMBER | | |
| | | | 5f. WORK UNIT NUMBER | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Children's Hospital Corporation Boston, MA 02115 | | | 8. PERFORMING ORGANIZATION REPORT | | |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | | |
| | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | | |
| 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT Our results show that neurons lacking Tsc1 results in a block in oligodendrocyte maturation and thus hypomyelination in the mouse brain. We show that Tsc-deficient neurons secrete excessive amounts of connective tissue growth factor (CTGF), which blocks the maturation of oligodendrocytes. We show here that CTGF is both necessary and sufficient to block the oligodendrocyte maturation. To determine whether CTGF is necessary <i>in vivo</i> , we generated mice lacking both Tsc1 and CTGF in neurons (Tsc1 ^{cc} ;CTGF ^{ff} ;SynCre ⁺) mice. In order to investigate the effect of CTGF on oligodendrocyte maturation, we propose to treat the oligodendrocytes with different domains of CTGF (Module I to IV). Moreover, in order to investigate the upstream regulation of CTGF expression in Tsc-deficient neurons, we tested Serum Response Factor (SRF) pathway, which was previously shown to affect CTGF expression. Our preliminary results show that SRF is downregulated in Tsc1 mutant brains and this can be rescued by rapamycin treatment suggesting a crosstalk between SRF and mTOR pathways. | | | | | |
| 15. SUBJECT TERMS Tuberous Sclerosis Complex, myelination, CTGF | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT | 18. NUMBER OF PAGES | 19a. NAME OF RESPONSIBLE PERSON USAMRMC |
| a. REPORT U | b. ABSTRACT U | c. THIS PAGE U | | | 19b. TELEPHONE NUMBER (include area code) |
| | | | UU | 11 | |

Table of Contents

| | <u>Page</u> |
|--|-------------|
| 1. Introduction..... | 4 |
| 2. Keywords..... | 4 |
| 3. Accomplishments..... | 4 |
| 4. Impact..... | 5 |
| 5. Changes/Problems..... | 6 |
| 6. Products..... | 6 |
| 7. Participants & Other Collaborating Organizations..... | 6 |
| 8. Special Reporting Requirements..... | 6 |
| 9. Appendices..... | 7 |
| 10. References..... | 11 |

INTRODUCTION

Tuberous sclerosis complex (TSC) is an autosomal dominant multisystem disorder caused by loss of either TSC1 or TSC2 function (Tsai and Sahin, 2011). TSC affects 1/6,000 individuals worldwide and affects multiple organs including the brain, skin, eyes, kidneys, heart, and lungs (Crino et al., 2006). TSC patients present with epilepsy (~90%), intellectual disability and autism (~50%), and other disorders including sleep disruption, attention-deficit hyperactivity disorder, and anxiety (Han and Sahin, 2011). The neuropathological findings in TSC are cortical tubers, subependymal nodules and subependymal giant cell astrocytomas (SEGAs) (DiMario, 2004). Another important yet not well-studied feature of TSC pathology in brain is hypomyelination (Ridler et al., 2001). Most recently using diffusion tensor imaging we observed abnormal white matter microstructure in patients with TSC that have autism compared to TSC patients without autism (Lewis et al., 2013; Peters et al., 2013). To uncover the underlying molecular mechanisms of hypomyelination in TSC, we investigated the role of neuronal factors affecting oligodendrocyte development in our *Tsc1^{cc}Syn/Cre⁺* mouse model, which lacks Tsc1 expression only in neurons. Here we show that, neurons lacking Tsc1 secrete excessive amounts of connective tissue growth factor (CTGF), which in turn blocks the maturation of oligodendrocytes, and thus myelination both *in vitro* and *in vivo*.

KEYWORDS

Tuberous Sclerosis Complex, myelination, CTGF

ACCOMPLISHMENTS:

We will review the accomplishments to date in each aim, as taken from the SOW (in **bold**). The figures are in the appendices.

Aim 1A: To determine the role of CTGF in hypomyelination in the mouse model.

I. We will generate two independent CTGF-shRNAs and confirm their efficacy in vitro.

We had previously shown that supernatant from HEK293T cells that express full length CTGF or commercially available recombinant CTGF reduce the number of MBP+ oligodendrocytes in cultures (Figure 1a-d). In addition, we had showed that adding neutralizing anti-CTGF antibody to conditioned medium from TSC-deficient neurons was able to ameliorate the effect of this conditioned medium on oligodendrocyte maturation and MBP expression (Figure 1e-f). We had proposed knocking down CTGF in TSC-deficient neurons as one way to verifying the role of CTGF in vitro. We did not proceed with these in vitro experiments since the *in vivo* experiments detailed below gave extremely compelling results allowing us to focus on the role of CTGF in the mouse model.

II. We will generate AAV2-CTGF-shRNA to reduce the expression of CTGF in vivo. AAV2-expressing scrambled shRNA as a control to the contralateral hemisphere. Alternatively, we will generate mice missing both Tsc1 and Ctgf genes under the Syn-Cre promoter (see grant proposal for details). Then we will follow look at MBP staining in the double knockout mice compared to single Tsc1 knockout mice.

We crossed *Tsc1;Syn/Cre⁺* mice with *CTGF^{ff}* mice (Kapoor et al., 2008) to generate the following groups of mice: wild type for both genes (*Tsc1^{ww};Syn/Cre⁺;CTGF^{+/+}*), mutant for *Tsc1* and wild type for *CTGF* (*Tsc1^{cc};Syn/Cre⁺;CTGF^{+/+}*) and mutant for both genes (*Tsc1^{cc};Syn/Cre⁺;CTGF^{ff}*). To our knowledge, this is the first mouse model lacking CTGF only in neurons. We stained the brain sections of control, mutant and double mutant with the MBP antibody to visualize myelination and found that loss of CTGF partially rescued the hypomyelination phenotype as assessed by the increased MBP

signal (Figure 1g-h). Interestingly, the number of mature oligodendrocytes, which were counted by CC1 staining, did not change between the mutant and double mutant (Figure 2a-c). We used CC1 expression to detect the mature oligodendrocytes by immunohistochemistry. This suggests that CTGF blocks the differentiation of oligodendrocytes to myelinating oligodendrocytes; however, another yet unknown factor affects the development of oligodendrocytes in mice lacking *Tsc1* in neurons. Moreover, the loss of CTGF in neurons, which are wild type for *Tsc1*, also enhanced myelination (Figure 2d-f). Together these data suggest that the neuronal CTGF is necessary for blocking myelination *in vivo*.

These results beg the question how TSC-deficiency regulates CTGF expression. We focused on the role of SRF. Previous reports show that the SRF functions as the repressor of *Ctgf* transcription (Stritt et al., 2009). Importantly, a previous study showed that neurons lacking SRF have higher expression of CTGF and thus blocks the maturation of oligodendrocytes (Stritt et al., 2009). We therefore analyzed the levels of SRF both in our mutant mice and in primary cortical neurons lacking *Tsc2*. Both SRF protein and transcript levels were decreased in *Tsc2* KD neurons compared to the control neurons (Figure 3a-c). In addition, we checked the transcript levels of other targets of SRF, *Egr1* and *Cyr61*. SRF functions as a transcription activator of *Egr1* (Kim et al., 2008), whereas it suppresses the transcription of *Cyr61* (Stritt et al., 2009), which is in the same CCN family of proteins as CTGF (Jun and Lau, 2011). In *Tsc2* KD cortical neurons, both the transcript levels of SRF and *Egr1* are decreased, whereas *Cyr61* is increased as *Ctgf* (Figure 3c). Moreover, when stained with SRF antibody, compared to the control mice, the mutant mice brain sections showed decreased intensity, suggesting SRF expression is diminished both *in vivo* and *in vitro*. Together, our data suggest that the upregulation of CTGF in *Tsc*-deficient neurons may be due to the downregulation of its repressor, SRF.

Aim 1B: To test whether CTGF expression is altered in human TSC brain.

We have initiated staining of CTGF in tuber specimens from TSC patients taken at the time of epilepsy surgery. We have so far stained paraffin sections of tubers from TSC patients and detected that CTGF is expressed in the tuber. We found that all phosphor-S6 positive cells are also CTGF-positive (Figure 4).

Aim 2: To examine the mechanisms by which CTGF regulates oligodendrocyte differentiation.

Our preliminary data from our first-year report suggests that the Mod-IV, which is one of the domains of CTGF is responsible for inhibiting oligodendrocyte maturation. To address this question systematically, we have been generating CTGF expression constructs, which express different domains of CTGF (Mod-I, Mod-II, Mod-III, Mod-IV and their combinations) to test the mutual and/or complementary effects of these different domains on oligodendrocyte maturation. So far we have generated the following FLAG-tagged constructs: Full-length, Module I, Module I+II. The cloning for the rest of the constructs is in progress. We had showed that the CM collected from HEK293T cells expressing the full-length FLAG-CTGF affects the maturation of oligodendrocytes. We will now test the individual modules for their effect on oligodendrocyte maturation assay.

IMPACT

In this study, we report that the double mutant, which lacks both *Tsc1* and CTGF in neurons rescue myelination. Moreover, we continue to work on the upstream regulators of CTGF expression and we focused on SRF pathway. In addition to our preliminary findings on SRF pathway in our previous report, here we show that SRF levels are decreased *in vivo* in mutant mice, and this can be rescued by rapamycin treatment. The crosstalk between SRF and mTOR pathways will not only provide a better understanding of regulation of CTGF expression but also help us to understand the basic control mechanisms of cell growth and survival. Better understanding of the role of CTGF in

myelination may provide insights and new therapeutic options for TSC and other diseases affecting the white matter.

CHANGES/PROBLEMS:

None

PRODUCTS:

We submitted the manuscript describing our findings for publication.

We have also published two review papers related to this topic and acknowledged the funding from the DOD:

[Autism and the synapse: emerging mechanisms and mechanism-based therapies.](#)

Ebrahimi-Fakhari D, **Sahin M.**

Curr Opin Neurol. 2015 Feb 19. [Epub ahead of print]

PMID: 25695134

Davis PE, Peters JM, Krueger D and **Sahin M.** Tuberous Sclerosis: A new frontier in targeted treatment of autism. Neurotherapeutics (in press).

PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS


None

SPECIAL REPORTING REQUIREMENTS

None

APPENDICES:

Figures 1-4






Figure 2. The number of CC1 positive mature oligodendrocyte number decreases in the *Tsc1* mutant mice, the double mutant, which lacks both *Tsc1* and *CTGF* does not rescue the number of CC1 positive cells (a-c). Mice, which are wild type for *Tsc1* but lacking *CTGF* in neurons, represent higher myelination as MBP intensity increases in the *CTGF* mutant compared to the wild type control (d-f).





Figure 3. SRF is downregulated in *Tsc*-deficient neurons. a) Immunoblotting of control and *Tsc2* KD cortical neurons showing Tsc2, SRF and Akt1 (loading control). b) Quantification of the SRF protein levels normalized to Akt1 (n=3). c) qRT-PCR of *Tsc2*, *Srf*, *Egr1*, *Cyr61* and *Ctgf* in *Tsc2* KD primary cortical neurons (n=3), showing the transcript levels. The dashed line represents the transcript levels in the control KD, which is set to 1. d) SRF staining of brain sections from control (n=3), *Tsc1* mutant (n=3) and rapamycin treated mutant (n=3) mice, scale bar corresponds to 200μm and corpus callosum (CC) is depicted. e) Quantifications of SRF intensity of the corresponding mice in arbitrary units (a.u.). (*p<0.05).



REFERENCES

- Crino, P.B., Nathanson, K.L., and Henske, E.P. (2006). The tuberous sclerosis complex. *N Engl J Med* 355,1345-1356.
- DiMario, F.J., Jr. (2004). Brain abnormalities in tuberous sclerosis complex. *J Child Neurol* 19,650-657.
- Han, J.M., and Sahin, M. (2011). TSC1/TSC2 signaling in the CNS. *FEBS Lett* 585,973-980.
- Heuer, H., Christ, S., Friedrichsen, S., Brauer, D., Winckler, M., Bauer, K., and Raivich, G. (2003). Connective tissue growth factor: a novel marker of layer VII neurons in the rat cerebral cortex. *Neuroscience* 119,43-52.
- Jun, J.I., and Lau, L.F. (2011). Taking aim at the extracellular matrix: CCN proteins as emerging therapeutic targets. *Nat Rev Drug Discov* 10,945-963.
- Kapoor, M., Liu, S., Huh, K., Parapuram, S., Kennedy, L., and Leask, A. (2008). Connective tissue growth factor promoter activity in normal and wounded skin. *Fibrogenesis Tissue Repair* 1,3.
- Kim, M.J., Kang, J.H., Chang, S.Y., Jang, H.J., Ryu, G.R., Ko, S.H., Jeong, I.K., Kim, M.S., and Jo, Y.H. (2008). Exendin-4 induction of Egr-1 expression in INS-1 beta-cells: interaction of SRF, not YY1, with SRE site of rat Egr-1 promoter. *J Cell Biochem* 104,2261-2271.
- Lewis, W.W., Sahin, M., Scherrer, B., Peters, J.M., Suarez, R.O., Vogel-Farley, V.K., Jeste, S.S., Gregas, M.C., Prabhu, S.P., Nelson, C.A., 3rd, and Warfield, S.K. (2013). Impaired language pathways in tuberous sclerosis complex patients with autism spectrum disorders. *Cereb Cortex* 23,1526-1532.
- Peters, J.M., Taquet, M., Prohl, A.K., Scherrer, B., van Eeghen, A.M., Prabhu, S.P., Sahin, M., and Warfield, S.K. (2013). Diffusion tensor imaging and related techniques in tuberous sclerosis complex: review and future directions. *Future Neurol* 8,583-597.
- Ridler, K., Bullmore, E.T., De Vries, P.J., Suckling, J., Barker, G.J., Meara, S.J., Williams, S.C., and Bolton, P.F. (2001). Widespread anatomical abnormalities of grey and white matter structure in tuberous sclerosis. *Psychol Med* 31,1437-1446.
- Stritt, C., Stern, S., Harting, K., Manke, T., Sinske, D., Schwarz, H., Vingron, M., Nordheim, A., and Knoll, B. (2009). Paracrine control of oligodendrocyte differentiation by SRF-directed neuronal gene expression. *Nat Neurosci* 12,418-427.
- Tsai, P., and Sahin, M. (2011). Mechanisms of neurocognitive dysfunction and therapeutic considerations in tuberous sclerosis complex. *Curr Opin Neurol* 24,106-113.